

Table 28. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 2003-04 season. There were n=466 licensed trappers in Mississippi in 2003-04.

Species	Harvest ^a	SE	Average Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	271.20	26.27	0.65	0.07	4.09	0.31	52	17.05	0.010
Raccoon	12,596.01	459.02	30.40	1.13	33.76	1.16	293	96.07	0.005
Muskrat	718.10	50.13	1.77	0.13	6.87	0.37	82	26.88	0.011
Red Fox	290.30	21.68	0.61	0.04	3.04	0.17	75	24.59	0.011
Gray Fox	1,685.75	144.21	3.37	0.15	7.97	0.65	166	54.42	0.013
Bobcat	2,010.42	104.56	4.81	0.25	8.09	0.38	195	63.93	0.012
Opossum	8,027.68	383.47	19.52	0.95	28.27	1.23	223	73.11	0.011
Otter	2,610.11	105.17	6.249	0.25	9.67	0.33	212	69.50	0.012
Spotted Skunk	61.11	13.62	0.06	0.014	4.80	0.85	10	3.27	0.004
Striped Skunk	1,030.04	77.26	2.35	0.16	7.42	0.48	109	35.74	0.012
Coyote	3,063.38	199.61	6.61	0.41	12.86	0.78	187	61.31	0.013
Weasel	20.37	5.94	0.052	0.02	4.00	0.82	4	1.31	0.003
Nutria	1,278.32	106.58	3.13	0.26	9.29	0.69	108	35.41	0.012
Beaver	14,085.68	596.59	34.68	1.49	42.88	1.68	258	84.59	0.009

^a n =366

^b n =305

^c Sample size for average catches per successful trapper

Table 29. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 2004-05 season. There were n=503 licensed trappers in Mississippi in 2004-05.

Species	Harvest ^a	SE	Average Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	316.66	31.70	0.67	0.07	4.26	0.35	57	18.10	0.010
Raccoon	14,199.98	666.66	31.46	1.37	36.44	1.64	299	94.92	0.006
Muskrat	616.37	47.81	1.43	0.11	5.99	0.36	79	25.08	0.012
Red Fox	385.72	35.13	0.92	0.08	3.75	0.29	79	25.08	0.012
Gray Fox	1,529.85	77.32	3.23	0.17	7.20	0.30	163	51.75	0.014
Bobcat	2,027.64	97.14	4.63	0.23	7.67	0.32	203	64.44	0.013
Opossum	6,889.54	325.33	14.67	0.63	23.09	0.98	229	72.70	0.012
Otter	2,736.53	115.96	6.29	0.26	10.34	0.37	203	64.44	0.013
Spotted Skunk	177.22	63.89	0.35	0.15	17.00	5.78	8	2.54	0.004
Striped Skunk	810.53	59.75	1.89	0.14	6.28	0.38	99	31.43	0.013
Coyote	3,287.74	273.06	7.50	0.66	14.50	1.14	174	55.24	0.014
Weasel	0.00	0.00	0.00	0.00	0.00	0.00	-	-	-
Nutria	3,332.05	945.66	8.09	2.30	22.43	6.33	114	36.19	0.013
Beaver	14,207.80	634.48	32.82	1.48	43.27	1.77	252	80.00	0.011

^a n =386

^b n =315

^c Sample size for average catches per successful trapper

Table 30. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 2005-06 season. There were n=476 licensed trappers in Mississippi in 2005-06.

Species	Harvest ^a	SE	Average Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	498.47	92.96	1.12	0.066	7.24	1.26	49	15.56	0.010
Raccoon	13,432.50	629.42	30.02	1.13	34.04	1.53	281	89.21	0.010
Muskrat	1,022.21	167.39	2.30	0.125	7.83	1.23	93	29.52	0.014
Red Fox	258.36	26.07	0.57	0.044	2.88	0.23	64	20.32	0.012
Gray Fox	1,272.14	76.98	2.79	0.149	6.29	0.32	144	45.71	0.015
Bobcat	2,071.09	104.61	4.62	0.25	7.34	0.33	201	63.81	0.015
Opossum	5,921.22	298.95	13.12	0.95	20.57	0.92	205	65.08	0.014
Otter	3,506.11	160.57	7.89	0.254	12.00	0.47	208	66.03	0.014
Spotted Skunk	241.51	113.09	0.55	0.014	15.64	7.21	11	3.49	0.006
Striped Skunk	751.21	58.69	1.68	0.16	6.08	0.37	88	27.94	0.014
Coyote	3,417.65	234.80	7.61	0.408	13.99	0.88	174	55.24	0.015
Weasel	2.81	1.064	0.01	0.0153	1.00	0.00	2	0.64	0.002
Nutria	1,954.55	149.24	4.38	0.26	10.96	0.73	127	40.32	0.015
Beaver	17,006.82	885.30	37.53	1.98	48.06	2.36	252	80.00	0.012

^a n =339

^b n =315

^c Sample size for average catches per successful trapper

Table 31. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 2006-07 season. There were n=488 licensed trappers in Mississippi in 2006-07.

Species	Harvest ^a	SE	Average Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	315.12	34.75	0.76	0.08	4.59	0.45	56	16.72	0.009
Raccoon	13,481.31	509.20	30.79	1.16	33.42	1.22	329	98.21	0.003
Muskrat	999.30	197.99	2.42	0.48	11.48	2.22	71	21.19	0.01
Red Fox	386.23	31.71	0.84	0.07	3.42	0.25	92	27.46	0.01
Gray Fox	1,842.87	140.37	3.67	0.17	8.40	0.61	179	53.43	0.01
Bobcat	2,247.50	83.06	5.18	0.19	7.97	0.26	230	68.66	0.01
Opossum	5,534.75	248.93	12.83	0.59	19.63	0.81	230	68.66	0.01
Otter	2,227.88	85.70	5.22	0.20	9.18	0.29	198	59.10	0.01
Spotted Skunk	50.27	11.14	0.12	0.03	3.73	0.71	11	3.28	0.004
Striped Skunk	1,050.79	114.53	2.47	0.28	7.58	0.79	113	33.73	0.01
Coyote	3,263.96	195.71	7.05	0.39	13.24	0.74	201	60.0	0.01
Weasel	1.23	0.53	0.003	0.001	1.00	-	1	0.30	0.003
Nutria	1,623.40	94.44	3.59	0.21	10.11	0.50	131	39.10	0.39
Beaver	13,471.50	567.48	31.76	1.35	41.46	1.63	265	79.10	0.79

^a n = 398

^b n = 335

^c Sample size for average catches per successful trapper

Table 32. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 2007-08 season. There were n=475 licensed trappers in Mississippi in 2007-08.

Species	Harvest ^a	SE	Average Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	177.68	20.09	0.49	0.06	3.85	0.37	39	12.87	0.008
Raccoon	11,538.59	469.79	31.67	1.26	36.08	1.38	270	89.11	0.007
Muskrat	852.87	190.22	2.37	0.53	13.85	3.03	52	17.16	0.009
Red Fox	305.61	25.11	0.82	0.07	3.69	0.26	70	23.10	0.01
Gray Fox	1,547.01	74.07	4.20	0.20	8.16	0.34	160	52.81	0.01
Bobcat	1,850.25	75.66	5.05	0.20	8.22	0.29	190	62.71	0.01
Opossum	6,021.01	306.62	16.65	0.84	26.20	1.22	194	64.03	0.01
Otter	1,395.39	54.16	3.86	0.15	6.62	0.21	178	58.75	0.01
Spotted Skunk	52.12	7.77	0.14	0.02	4.00	0.38	11	3.63	0.004
Striped Skunk	868.27	58.78	2.41	0.16	8.06	0.46	91	30.03	0.01
Coyote	2,700.75	158.20	7.47	0.43	13.49	0.73	169	55.78	0.01
Weasel	18.95	6.16	0.05	0.02	5.33	1.52	3	0.99	0.002
Nutria	1,265.09	72.32	3.45	0.20	9.71	0.46	110	36.30	0.01
Beaver	11,925.94	524.87	32.67	1.41	45.76	1.84	220	72.61	0.01

^a n = 401

^b n = 303

^c Sample size for average catches per successful trapper

Table 33. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 20008-09 season. There were n=538 licensed trappers in Mississippi in 2008-09.

Species	Harvest ^a	SE	Average Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	105.86	13.36	0.26	0.03	3.27	0.31	26	8.08	0.007
Raccoon	12,916.98	493.48	30.61	1.17	34.23	1.22	303	94.10	0.006
Muskrat	706.13	144.33	1.76	0.36	12.60	2.48	45	13.98	0.009
Red Fox	328.78	30.91	0.63	0.06	3.52	0.29	75	23.29	0.010
Gray Fox	1,555.47	86.20	3.45	0.18	7.66	0.37	163	50.62	0.012
Bobcat	1,753.48	95.72	4.10	0.22	7.91	0.38	178	55.28	0.012
Opossum	7,757.41	513.85	18.01	1.25	24.72	1.58	252	78.26	0.010
Otter	1,689.97	84.55	4.09	0.20	8.03	0.34	169	52.48	0.012
Spotted Skunk	33.63	7.03	0.05	0.01	3.86	0.53	7	2.17	0.004
Striped Skunk	780.85	49.22	1.63	0.10	6.21	0.31	101	31.37	0.012
Coyote	3,442.20	230.48	6.75	0.39	16.26	1.00	170	52.80	0.012
Weasel	0.00	0.00	0.00	0.00	0.00	0.00	0	0.00	0.00
Nutria	1,560.45	122.63	3.57	0.29	10.53	0.75	119	36.96	0.012
Beaver	10,544.55	466.70	25.39	0.12	36.03	1.43	235	72.98	0.011

^a n = 432

^b n = 322

^c Sample size for average catches per successful trapper

Table 34. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 2009-10 season. There were n=426 licensed trappers in Mississippi in 2009-10.

Species	Harvest ^a	SE	Average Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	143.67	24.34	0.29	0.04	4.42	0.66	26	10.36	0.009
Raccoon	8,390.08	313.62	26.13	0.93	29.85	0.99	225	89.64	0.009
Muskrat	391.02	65.09	1.20	0.21	7.11	1.11	44	17.53	0.011
Red Fox	173.65	12.89	0.54	0.04	2.53	0.13	55	21.91	0.012
Gray Fox	1,065.62	62.55	3.33	0.19	6.77	0.34	126	50.20	0.014
Bobcat	1,119.34	56.76	3.51	0.17	6.64	0.27	135	53.78	0.014
Opossum	5,234.43	255.11	16.00	0.76	24.94	1.05	168	66.93	0.013
Otter	1,469.14	79.51	4.63	0.24	9.12	0.41	129	51.39	0.014
Spotted Skunk	14.99	3.04	0.05	0.01	2.00	0.20	6	2.39	0.004
Striped Skunk	454.73	40.06	1.29	0.11	5.97	0.43	61	24.30	0.012
Coyote	2,570.99	188.98	7.57	0.55	15.36	1.03	134	53.39	0.014
Weasel	2.50	1.12	0.01	0.00	2.00		1	0.40	0.002
Nutria	1,173.06	101.16	3.31	0.28	10.67	0.81	88	35.06	0.013
Beaver	9,359.51	469.96	29.29	1.44	40.50	1.83	185	73.71	0.012

^a n = 341

^b n = 251

^c Sample size for average catches per successful trapper

Table 35. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 2010-11 season. There were n=478 licensed trappers in Mississippi in 2010-11.

Species	Harvest ^a	SE	Average Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	93	10.54	0.23	0.03	2.37	0.18	30	10.20	0.86
Raccoon	12,569	502.47	31.70	1.25	35.68	1.32	269	91.50	0.79
Muskrat	466	62.18	1.19	0.16	6.36	0.76	56	19.05	1.12
Red Fox	229	19.58	0.54	0.05	3.07	0.19	57	19.39	1.42
Gray Fox	1,435	76.26	3.59	0.19	7.46	0.32	147	50.00	1.42
Bobcat	1,714	84.02	4.34	0.21	8.29	0.33	158	53.74	1.28
Opossum	7,647	510.25	19.13	1.30	27.81	1.75	210	71.43	1.42
Otter	2,010	105.33	5.01	0.26	9.97	0.43	154	52.38	1.42
Spotted Skunk	14	3.97	0.04	0.01	2.75	0.42	4	1.31	0.33
Striped Skunk	665	50.77	1.68	0.13	5.64	0.35	90	30.61	1.31
Coyote	3,374	207.43	8.37	0.52	15.61	0.86	165	56.12	1.41
Weasel	1	0.64	0.00	0.00	1.0	-	1	0.34	0.17
Nutria	939	68.23	2.37	0.17	8.15	0.47	88	29.93	1.30
Beaver	10,925	578.69	26.13	1.38	38.44	1.87	217	73.81	1.25

^a n = 365

^b n = 294

^c Sample size for average catches per successful trapper

Table 36. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 2011-12 season. There were n=504 licensed trappers in Mississippi in 2011-12.

Species	Harvest ^a	SE	Average Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	171	21.26	0.43	0.05	3.94	0.37	33	10.86	0.88
Raccoon	10,608	418.42	25.93	0.99	29.21	1.06	276	90.79	0.81
Muskrat	251	24.63	0.63	0.06	4.24	0.30	45	14.80	0.10
Red Fox	238	19.98	0.59	0.05	2.66	0.17	68	22.37	1.17
Gray Fox	1,609	92.81	3.99	0.23	8.15	0.40	150	49.34	1.41
Bobcat	1,880	107.85	4.63	0.26	8.56	0.43	167	54.93	1.40
Opossum	4,947	214.30	11.72	0.49	18.70	0.68	201	66.12	1.33
Otter	2,362	123.81	5.65	0.30	10.95	0.48	164	53.95	1.40
Spotted Skunk	26	7.67	0.07	0.02	4.00	0.86	5	1.65	0.36
Striped Skunk	598	40.64	1.50	0.10	4.79	0.25	95	31.25	1.31
Coyote	3,574	249.42	8.85	0.61	16.87	1.07	161	52.96	1.41
Weasel	1	0.65	0.00	0.00	1.00	-	1	0.33	0.16
Nutria	707	46.76	1.60	0.10	5.77	0.29	93	30.59	1.30
Beaver	11,299	600.57	27.99	1.46	36.23	1.79	237	77.96	1.17

^a n = 383

^b n = 304

^c Sample size for average catches per successful trapper

Table 1. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 2013-14 season. There were n=511 licensed trappers in Mississippi in 2012-13.

Species	Harvest ^a	SE	Average Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	133		0.22		4.10			5.47	
Raccoon	12,397		21.05		28.38			74.17	
Muskrat	643		1.09		8.33			13.11	
Red Fox	316		0.54		3.11			17.22	
Gray Fox	2,139		3.63		9.57			37.97	
Bobcat	1,939		3.29		7.47			44.03	
Opossum	7,818		13.27		24.14			55.00	
Otter	2,359		4.01		9.48			42.27	
Spotted Skunk	73		0.12		3.15			3.91	
Striped Skunk	816		1.39		5.71			24.27	
Coyote	3,818		6.48		14.99			43.25	
Weasel	6		0.01		1.00			0.98	
Nutria	1,208		2.05		8.96			22.90	
Beaver	11,168		18.96		33.30			56.95	

^a n = 511

^b n =

^c Sample size for average catches per successful trapper

Table 1. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trappers, and percent successful trappers by furbearer species for the 2013-14 season. There were n=707 licensed trappers in Mississippi in 2013-14.

Species	Harvest ^a	SE	Average Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	135	16.89	0.19	0.02	3.27	0.33	33	5.84	0.44
Raccoon	18,423	462.31	26.06	0.65	29.86	0.72	493	87.26	0.63
Muskrat	616	49.88	0.87	0.07	6.15	0.41	80	14.16	0.66
Red Fox	564	32.03	0.80	0.05	2.91	0.14	155	27.43	0.84
Gray Fox	2,632	107.88	3.72	0.15	7.82	0.28	269	47.61	0.94
Bobcat	2,484	87.69	3.51	0.12	6.59	0.20	301	53.27	0.94
Opossum	8,540	339.56	12.08	0.48	18.80	0.70	363	64.25	0.90
Otter	2,519	94.21	3.56	0.13	7.89	0.25	255	45.13	0.94
Spotted Skunk	43	8.29	0.06	0.01	3.78	0.51	9	1.59	0.24
Striped Skunk	1,042	80.82	1.47	0.11	5.95	0.42	140	24.78	0.81
Coyote	4,204	238.62	5.95	0.34	12.73	0.68	264	46.73	0.94
Weasel	4	0.97	0.01	0.00	1.00	0.00	3	0.53	0.14
Nutria	1,283	108.26	1.81	0.15	8.27	0.63	124	21.95	0.78
Beaver	11,789	469.99	16.67	0.66	26.24	0.98	359	63.54	0.91

^a n = 383

^b n = 304

^c Sample size for average catches per successful trapper